

Induction treatment consisted of weekly injections until the lesions were inactive. Maintenance therapy consisted of one injection each 2-4 weeks continuously until relapse, then the weekly schedule was re-started.

The study protocol was approved by the research ethics committee of the Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand (No. 55/2003).

Three hundred and sixty-three patients (568 eyes) were treated and followed for a mean period of 25.3 weeks (range, 18 to 118 weeks). Table 1 shows the

Characteristics	Number (%)
No. of patients	363
Age (yrs)	
Mean \pm SD	34.5 \pm 7.6
Range	5-64
Sex	
Male	163 (45%)
Female	200 (55%)
Laterality at first	
Unilateral	203 (56%)
Bilateral	160 (44%)
Visual Acuity at initial visit (eyes) (N = 523 eyes)	
6/6-6/18	216 (41%)
< 6/18-3/60	142 (27%)
< 3/60 - PL	153 (29%)
No PL	12 (2%)
Location of lesions at initial visit (eyes)	
Zone 1	13 (2%)
Zone 2	168 (32%)
Zone 3	6 (1%)
Zone 1 and zone 2	229 (44%)
Zone 2 and zone 3	58 (11%)
Zone 1 and zone 2 and zone 3	49 (9%)

18/11/05, 5:07 PM

Table 2. Results of Intravitreal Ganciclovir Treatment in CMV Retinitis Patients

Results	Number (%)
No.of patients who developed CMV retinitis in the other eye	45 (22%)
Time of developed CMVR in the other eye (wks)	
Mean \pm SD	9.5 \pm 10.0
Range	1-43
Visual outcome (eyes) (N = 568 eyes)	
Stable	343 (60%)
Improved	76 (13%)
Decreased	149 (26%)
Complications (eyes)	
Endophthalmitis	6
Retinal detachment	33
Vitreous hemorrhage	6
Cataract	2

CMV = Cytomegalovirus

baseline ocular characteristics of the patients at their first visit. The patients ranged in age from 5 to 64 years (mean 34.5 years). One hundred and sixty-three (45%) were male. Retinitis was initially unilateral in 203 patients (56%) and bilateral in 160 (44%). The visual acuity at initial presentation was in the 6/6 to 6/18 range in 216 eyes (41%), <6/18-3/60 in 142 (27%), <3/60-PL in 153 (29%), and no PL in 12 (2%). Retinitis affected only zone 1 in 13 eyes (2%), zone 2 in 168 (32%), zone 3 in 6 (1%), both zone 1 and zone 2 in 229 (44%), zone 2 and zone 3 in 58 (11%), and all three zones in 49 (9%) at the initial visit.

The results of intravitreal ganciclovir treatment are summarized in Table 2. Involvement of the fellow eye developed in 45 (22%) of 203 patients during treatment of the first eye, between 1 and 43 weeks (mean 9.5 weeks). In total, 568 eyes were injected with intravitreal ganciclovir. At the time of last follow up, 343 (60%) had stable vision, 76 (13%) had improved vision, and 149 (26%) had decreased vision. Complications in 568 treated eyes included 6 with endophthalmitis, 33 with retinal detachment, 6 with vitreous hemorrhages, and 2 with cataract.

Discussion

CMV retinitis is the most common sight-threatening complication in acquired immune deficiency syndrome (AIDS), occurring in 33 % of patients in Chiang Mai study⁽¹⁹⁾. Systemic treatment with ganciclovir has been the mainstay of management⁽¹⁻⁸⁾, but frequently was complicated by its toxicity and deterioration in quality of life^(5,6,9,10). Intravitreal ganciclovir has shown to be effective in stopping the progression

of CMV retinitis in AIDS patients⁽¹¹⁻¹⁶⁾. The authors therefore evaluated the efficacy of this drug in terms of the visual outcome, and complications of the treatment.

In the present study, eyes with stabilized, improved, and decreased visual outcome were 60%, 13%, and 26%, respectively (Table 2). If the eyes with initial visual acuity of no PL (12 eyes or 2%) were not taken into account for stabilized visual outcome, therefore 71% of eyes could preserve vision no worse than before by this treatment modality. Nevertheless, the initial visual acuity, location and extent of lesions, complications of diseases, and other AIDS-related disorders such as retinal ischemic associated with microvascular diseases and lesions involving intracranial visual pathways should also be considered to be the other factors that affected the visual outcome⁽⁴⁾.

Bilateral disease occurred in 45 (22%) of 203 patients, who first came with unilateral involvement. It is known that intravitreal therapy is purely palliative as it has no effect on the systemic cytomegalovirus infection. However, systemic ganciclovir may not offer the expected advantage in this respect, since contralateral retinitis has been reported in 15-68% of patients receiving intravenous maintenance therapy^(5-6,9-10). In addition, frequent follow up meant that all second eye infections might be recognized early.

Six eyes had endophthalmitis, representing 1.1 % of 568 treated eyes. Multiple injections that required to maintain remission may be the risk factor that attributes to this serious complication of intravitreal injections. A previous author's study using intravitreal foscarnet reported an incidence of endophthalmitis of 1% of treated eyes⁽¹⁸⁾. Some studies advocated intra-

There were 33 eyes (5.8%) with retinal detachments and 6 (1.2 %) with intravitreal hemorrhages. These rates are not different from those reported in other intravitreal series^(11,12,14,16,18). It was accepted that intravitreal injection was not attributed to these complications, since retinal detachment was a frequent complications of this disease, with an incidence varying from 11% to 29%^(20,21). Risk factor for development of rhegmatogenous retinal detachment in patients with CMV retinitis were peripheral involvement greater than 25%, the presence of active retinitis, greater patient age, and lower CD4+ cell counts^(20,21). It was also reported that intravitreal therapy offered a significant benefit over systemic therapy in the risk of CMV-related retinal detachment⁽²²⁾. Although vitreous anomalies induced by ganciclovir (pH 10.14) was suspected to be a contributing factor⁽¹⁴⁾, a previous author's study using foscarnet (pH 7.4) did not demonstrated this advantage⁽¹⁸⁾.

Although there was a previous study using intravitreal foscarnet⁽¹⁸⁾, the authors did not compare the result of both studies with statistical analysis. They were both not randomized controlled trials, and also shared the weakness inherent in all retrospective studies.

Acknowledgement

References

- Cytomegalovirus retinitis and response to therapy with ganciclovir. *Ophthalmology* 1987; 94: 425-34.
2. Holland GN, Sidikaro Y, Kreiger AE, Hardy D, Sakamoto MJ, Frenkel LM, et al. Treatment of cytomegalovirus retinopathy with ganciclovir. *Ophthalmology* 1987; 94: 815-23.
3. Jabs DA, Newman C, de Bustros S, Polk BF. Treatment of cytomegalovirus retinitis with ganciclovir. *Ophthalmology* 1987; 94: 824-30.
4. Holland GN, Buhles WC, Mastre B, Kaplan H, the UCLA CMV Retinopathy Study Group. A controlled retrospective study of ganciclovir treatment for cytomegalovirus retinopathy: use of a standard system for assessment of disease outcome. *Arch Ophthalmol* 1989; 107: 1759-66.
5. Jacobson MA, O'Donnell JJ. Approaches to the treatment of cytomegalovirus retinitis : ganciclovir and foscarnet. *J Acquir Immune Defic Syndr* 1991; 4(Suppl 1): S11-5.
6. Studies of Ocular Complications of AIDS Research Group in collaboration with the AIDS Clinical Trials Group. Mortality in patients with the acquired immunodeficiency syndrome treated with either foscarnet or ganciclovir for cytomegalovirus retinitis. *N Engl J Med* 1992; 326: 213-20.
7. Studies of Ocular Complications of AIDS (SOCA) Research Group in collaboration with the AIDS Clinical Trials Group (ACTG). Foscarnet-ganciclovir CMV retinitis trial: IV. Visual outcomes. *Ophthalmology* 1994; 101: 1250-61.
8. Studies of Ocular Complications of AIDS (SOCA) Research Group in collaboration with the AIDS Clinical Trials Group (ACTG). Combination foscarnet and ganciclovir therapy vs monotherapy for the treatment of relapse CMV retinitis in patients with AIDS. *Arch Ophthalmol* 1996; 114: 23-33.
9. Gross JG, Bozzette SA, Mattheus WC, Spector SA, Abraham IA, McCutchan JA, et al. Longitudinal study of cytomegalovirus retinitis in acquired immune deficiency syndrome. *Ophthalmology* 1990; 97: 681-6.
10. Peter BS, Beck EJ, Anderson S, Coleman D, Coker R, Main J, et al. Cytomegalovirus infection in AIDS. Patterns of disease, response to therapy and trends in survival. *J Infect* 1991; 23: 129-37.
11. Ussery FM, Gibson SR, Conklin RH, Pitot DF, Stool EW, Conklin AJ. Intravitreal ganciclovir in the treatment of AIDS associated cytomegalovirus retinitis. *Ophthalmology* 1988; 95: 640-8.
12. Heinemann MH. Long-term intravitreal ganciclovir therapy for cytomegalovirus retinopathy. *Arch*

- Ophthalmol 1989; 107: 1767-72.
13. Cantrill HL, Henry K, Melroe NH, Knobloch WH, Ramsay RC, Balfour HH. Treatment of cytomegalovirus retinitis with intravitreal ganciclovir: long-term results. *Ophthalmology* 1989; 96: 367-74.
 14. Cochereau-Massin I, Lehoang P, Lautier-Frau M, Zazoun L, Marcel P, Robinet M, et al. Efficacy and tolerance of intravitreal ganciclovir in cytomegalovirus retinitis in acquired immune deficiency syndrome. *Ophthalmology* 1991; 98: 1348-55.
 15. Morlet N, Young S, Naidoo D, Graham G, Coroneo MT. High dose intravitreal ganciclovir injection provides a prolonged therapeutic intraocular concentration. *Br J Ophthalmol* 1996; 80: 214-6.
 16. Young S, Morlet N, Besen G, Wiley CA, Jones P, Gold J, et al. High-dose (2000- μ g) intravitreal ganciclovir in the treatment of cytomegalovirus retinitis. *Ophthalmology* 1998; 105: 1404-10.
 17. Holland GN, Tufail A, Jordan MC. Cytomegalovirus diseases. In: Pepose JS, Holland GN, Wilhelmus KR, editors. *Ocular infection and immunity*. St. Louis: Mosby, 1996 : 1088-129.
 18. Ausayakhun S, Watananikorn S, Ngamtiphakorn S, Prasitsilp J. Intravitreal foscarnet for cytomegalovirus retinitis in patients with AIDS. *J Med Assoc Thai* 2005; 88: 103-7.
 19. Ausayakhun S, Watananikorn S, Ittipunkul N, Chaidaroon W, Patikulsila P, Patikulsila D. Epidemiology of the ocular complications of HIV infections in Chiang Mai. *J Med Assoc Thai* 2003; 86: 399-406.
 20. Freeman WR, Friedberg DN, Berry C, Quiceno JJ, Behette M, Fullerton SC, et al. Risk factors for development of rhegmatogenous retinal detachment in patients with cytomegalovirus retinitis. *Am J Ophthalmol* 1993; 116: 713-20.
 21. Cassoux N, Bodaghi B, Lautier-Frau M, Fardeau C, Lehoang P. Current status of retinal detachment in AIDS patients. *J Fr Ophthalmol* 2000; 23: 1031-4.
 22. Young S, McCluskey P, Minassian DC, Joblin P, Jones C, Coroneo M, et al. Retinal detachment in cytomegalovirus retinitis: intravenous versus intravitreal therapy. *Clin Experiment Ophthalmol* 2003; 31: 96-102.

การรักษาจอประสาทตาอักเสบจากไซโตเมกะโลไวรัสในผู้ป่วยเอดส์ โดยการฉีดยาแกนไซโคล- เวียร์เข้าวันตา

สมสงวน อัญญคุณ, ประภา ยวเวส, สุภพ งามทิพากร, จิระเดช ประสิทธิศิลป์

ความเป็นมา: โรคจอประสาทตาอักเสบจากไซโตเมกะโลไวรัส เป็นโรคติดเชื้อฉวยโอกาสทางตาที่พบบ่อยที่สุดในผู้ป่วยกลุ่มอาการภูมิคุ้มกันบกพร่อง และมักทำให้ตาบอดได้ถ้าไม่ได้รับการรักษา การฉีดยาแกนไซโคลเวียร์เข้าวันตาสามารถยับยั้งโรคได้

วัตถุประสงค์: เพื่อศึกษาประสิทธิภาพและผลแทรกซ้อนของการฉีดยาแกนไซโคลเวียร์เข้าวันตา (2 มิลลิกรัม ใน 0.1 มิลลิลิตร) ในการยับยั้งโรคจอประสาทตาอักเสบจากไซโตเมกะโลไวรัส

รูปแบบการวิจัย: การศึกษาแบบย้อนหลังในกลุ่มผู้ป่วย

วัสดุและวิธีการ: ผู้ป่วยโรคจอประสาทตาอักเสบจากไซโตเมกะโลไวรัส 363 คน ที่มารักษาที่โรงพยาบาลมหาราชนครเชียงใหม่ ตั้งแต่มิถุนายน พ.ศ. 2544 ถึง ธันวาคม พ.ศ. 2546 โดยในระยะเริ่มต้นทำการฉีดยาสัปดาห์ละครั้งจนรอยโรคสงบ แล้วตามด้วยการฉีด 2-4 สัปดาห์ต่อครั้งไปตลอดจนรอยโรคลุกลามใหม่ก็จะเริ่มต้นฉีดยาสัปดาห์ละครั้งใหม่

ผลการศึกษา: ตาที่ได้รับการรักษา 568 ตา พบว่า มีสภาพการมองเห็นเมื่อสิ้นสุดการรักษา (ระยะเวลาเฉลี่ย 25.7 สัปดาห์) ดังนี้ สภาพการมองเห็นเท่าเดิม 343 ตา (60%) สภาพการมองเห็นดีขึ้น 76 ตา (13%) และสภาพการมองเห็นลดลง 149 ตา (26%) ภาวะแทรกซ้อนที่พบคือ จอภาพตาหลุด 33 ตา เลือดออกในน้ำวันตา 6 ตา การอักเสบในลูกตา 6 ตา และต้อกระจก 2 ตา และผู้ป่วย 22% เกิดรอยโรคในตาอีกข้างหนึ่งในระหว่างการรักษาตาข้างแรก

สรุป: การรักษาด้วยการฉีดยาแกนไซโคลเวียร์เข้าวันตาเป็นอีกทางเลือกหนึ่งในผู้ป่วยที่ไม่สามารถให้ยาต้านเชื้อไซโตเมกะโลไวรัสทางเส้นเลือดได้ แต่ผลแทรกซ้อนจากการฉีดยาเข้าวันตาเป็นสิ่งที่ต้องคำนึงในการรักษาด้วย
